## This Page Is Inserted by IFW Operations and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

### IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

#### WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



#### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

(11) International Publication Number:

WO 98/37866

A61K 7/50, 7/48

A1 (43) International Publication Date:

3 September 1998 (03.09.98)

(21) International Application Number:

PCT/EP98/00155

(22) International Filing Date:

12 January 1998 (12.01.98)

(30) Priority Data:

08/810.114

25 February 1997 (25.02.97)

US

(71) Applicant (for AU BB CA GB GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BO

(71) Applicant (for all designated States except AU BB CA GB GH GM IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

- (72) Inventors: FUJIWARA, Mitsuko; 2310 Shurts Circle, Urbana, IL 61801 (US). VINCENT, Carol, Kregler, 18 Wolfe Drive, Wanaque, NJ 07465 (US). ANANTHAPADMANABHAN, Kavssery, Parameswaran; 23 Vanderbilt Drive, Highland Mills, NY 10930 (US). VILLA, Virgilio, Barba; 140 Grove Street, Bergenfield, NJ 07621 (US).
- (74) Agent: MOLE, Peter, Geoffrey; Unilever plc, Patent Division. Colworth House, Shambrook, Bedford MK44 1LQ (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

(54) Title: MILD ANTIMICROBIAL LIQUID CLEANSING FORMULATIONS COMPRISING HYDROXY ACID BUFFERING COMPOUND OR COMPOUNDS AS POTENTIATOR OF ANTIMICROBIAL EFFECTIVENESS

#### (57) Abstract

The present invention relates to liquid skin cleansing compositions comprising (1) mild surfactant systems; (2) 0.5 % to 9 % by wt. of a hydroxy carboxylic compound or compounds which buffer the pH of the composition; and (3) 1 % to 99 % water to potentiate the bactericidal activity. In a second embodiment of the invention, the buffering compound or compounds potentiates antibacterial effect in compositions already containing an antibacterial agent.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

·							= =
AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegai
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HŲ	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# MILD ANTIMICROBIAL LIQUID CLEANSING FORMULATIONS COMPRISING HYDROXY ACID BUFFERING COMPOUND OR COMPOUNDS AS POTENTIATOR OF ANTIMICROBIAL EFFECTIVENESS

#### 5

#### RELATED APPLICATIONS

The subject application is a continuation-in-part application of U.S. Serial No. 08/252,298, now allowed.

#### 10

15

#### BACKGROUND OF THE INVENTION

The present invention relates to one-phase liquid cleansing compositions having enhanced antimicrobial effectiveness. More specifically, the invention relates to a hydroxy acid compound or compounds which potentiate the antibacterial activity of liquid cleaning formulations by buffering the pH of the formulation such that the pH will rise no higher than 5.0, preferably between 2.5 to 5.0 under in use conditions (as opposed to initial pH).

20

25

30

There is a large demand in the market for mild liquid cleansing formulations which additionally have an antibacterial effect. Antibacterial cleansers are preferred because they kill germs and mild personal cleansers are preferred to minimize skin irritation, dryness, etc. However, the combination of mild cleansing formulations and strong antibacterial effect is difficult to achieve. Thus, for example, while soaps provide antibacterial effects, they are not mild to the skin. When very mild non-soap surfactants are used, antibacterial effect is greatly compromised.

The balancing act between providing mildness and effective antibacterial effectiveness is recognized for example in International Publication WO 92/18100. In this

publication, improved clinical mildness is said to be provided through the use of a water soluble cationic polymer (see page 10, lines 24-29). Cationic polymer is apparently used instead of additional ethoxylated surfactant because the percent of ethoxylated mildness surfactant must be minimized in order not to affect antibacterial effectiveness (page 7, lines 4-6).

Another approach to providing mildness effect without

affecting antibacterial properties is that which appears to
be used by Dial in, for example, Liquid Dial Plus with

Moisturizers Antibacterial Soap<sup>(R)</sup>. Here, mildness benefits
are apparently provided by the use of moisturizing compounds
rather than by the use of very mild surfactants alone

(which, as indicated above, compromises antibacterial
effectiveness).

In both cases, it can be readily seen that it is extremely difficult to provide effective antibacterial action in the presence of very mild surfactants, to use larger amounts of harsher surfactants or soaps and to mask the effects of the harshness by providing cationic mildness conditioning agents (WO 92/18100) or moisturizers (as in the Dial product).

25

30

20

It would be greatly beneficial if antibacterial effectiveness could be provided either by providing a compound or compounds which alone or together buffer pH of a liquid composition at a pH low enough to provide antibacterial effectiveness for that composition formulation (while maintaining stability of composition); or by providing a compound or compounds which alone or together buffer pH of a liquid composition containing anti-bacterial agent thereby enhancing (i.e., potentiating) the effect of

the antibacterial agent even in compositions with very mild surfactant systems.

Fatty acids and their ester derivatives have been used to provide antimicrobial effectiveness in foods, pharmaceuticals and cosmetics (see, for example EP 0,244,144; U.S. 4,002,775; U.S. 4,406,884; U.S. 4,997,851 and Kabara in JAOCS, vol. 61, No. 2, (February, 1984)).

The use of short chain fatty acids generally as potentiators of germicides is also known. These fatty acids, for example, have been used as potentiators with halogenated germicides at high pH and with isethiazolones (see FR 2,223,049 and EP 488,606).

15

20

25

35

5

U.S. 3,218,260 to Lewandowski discloses cleaner compositions containing various organic or inorganic acids. The pH of these compositions (less than 2) is well below the pH of the skin cleansing compositions of the present invention.

In none of these references is it taught or suggested that one or more compounds be used either to enhance antibacterial effect in liquid skin cleansing compositions or to potentiate antibacterial compounds which may already be present in liquid skin cleansing compositions at the pH specified by the claims of the subject invention.

Further, none of these references relate to use of hydroxy carboxylic acid (e.g., lactic acid).

U.S. Patent No. 5,132,037 to Greene et al. teaches aqueous compositions in which  $C_8$ - $C_{22}$  free fatty acids may be used. All examples (palmitic, stearic) are clearly directed to longer chain fatty acids and there is absolutely no

recognition of the antibacterial or potentiating effect of lower chain fatty acids. Also, there is no teaching or suggestion of hydroxy carboxylic acids.

5 U.S. Patent No. 5,137,715 to Hoshowski et al. teaches shampoo conditioner compositions wherein the pH of the composition can be in the range of 2.5 to 7.0. The invention requires a polymeric amidoamine compound (substantive compound which imparts conditioning and does not adversely affect foam of anionic; see column 11, line 63 to column 12, line 36). It is further taught that an acid is required to neutralize the amidoamine and one acid which is said to be used for this purpose is citric acid (see column 13, lines 49-65).

15

20

The compositions of Hoshowski, while stable, were only stable when using the specific amidoamine of formula I (Example 13 of the patent notes that an extremely similar amidoamine, represented by Formula V, caused instability at pH below 6.0) and, according to examples, 2% citric acid was used.

In general Hoshowski et al. makes clear that most amidoamines would cause instability. More specifically, 25 applicants tried the amidoamine of Formula I in compositions of the subject invention and also found instability. Applicants are not certain whether this instability was due to large amounts of hydroxy acid (applicants use minimum 0.5% lactic acid versus 0.2% citric acid exemplified); whether it was due to the specific hydroxy acid used; or 30 whether it was due to the specific surfactant system. is clear, however, is that there is no such instability in the system of the invention without the amidoamine of Formula I while there is such instability using the 35 amidoamine in the same system.

- U.S. Patent No. 5,002,180 to Schmidt et al. teach skin cleansing aerosol mousse emulsions comprising:
  - (A) 88% to 97% of a concentrate comprising:
    - (1) 3%-20% anionic or amphoteric;
    - (2) 0.05 to 5% polymeric skin feel aid;
    - (3) 10% to 60% moisturizers (which can be lactic acid); and
    - (4) water; and
  - (B) 3% to 12% propellant.

15

20

30

5

This reference differs from the subject compositions in a number of ways. First the lactic acid, if used, is used as moisturizing component and must comprise 10% or greater of composition whereas upper level of the hydroxy carboxylic acid of invention (to provide bactericidal effect) is about 9%. Further, the reference is not a single phase composition but comprises propellant (to form mousse). While not wishing to be bound by theory, bactericidal effect of hydroxy acid of invention are believed to be due largely to single phase systems of invention. In a multiphase, it is believed surfactant would not have time to solubilize and enter liquid phase and therefore could not deliver antibacterial activity.

- In short, applicants have now found that one or more hydroxy compounds may be used to buffer low pH within a defined low pH range and to therefore:
  - (1) enhance the antibacterial effect of liquid skin cleansing compositions; and/or
  - (2) potentiate antimicrobial effect of liquid skin cleansing compositions which already contain an antimicrobial agent.

The single phase compositions of the invention are free of amidoamines generally and more specifically, of the amidoamines described in U.S. Patent No. 5,137,715 to Hoshowski.

5

#### BRIEF SUMMARY OF THE INVENTION

The present invention relates to liquid skin cleansing compositions comprising:

10

15

20

25

- (1) any mild surfactant system (i.e., any one or more surfactants which alone or together are demonstrated by clinical tests to be milder than soap itself) in an amount of from about 1-99% by wt., preferably 2-85% by wt., more preferably 3-40% by wt. surfactant system;
- (2) 0.5 to about 9%, preferably 0.5 to 5% by weight of a hydroxy carboxylic compound or compounds (e.g., lactic acid) which alone or together buffer the pH of the liquid skin cleanser composition such that the pH is no higher than 5.5 under in-use conditions (i.e., 1:0.5 to 1:100 dilution, preferably 1:1 to 1:25 dilution of product in H<sub>2</sub>O); and
- (3) 1% to 99% by wt., preferably 15 to 97%, most preferably 60 to 97% by wt. water.

More specifically, the composition may comprise:

- 30 (1) 1% to 99% by wt. of surfactant system comprising:
  - (a) 1 to 30% by wt. of at least one anionic surfactant;
  - (b) 0.5% to 15% amphoteric surfactant;
  - (2) 0.5 to 9% hydroxy acid; and
- 35 (3) 1% to 99% water.

30

In a second embodiment of the invention, the liquid skin cleansing composition comprises 0.0001% to 5% by weight of an antibacterial agent and the buffering compound or compounds act to potentiate the antimicrobial effect of the composition.

#### BRIEF DESCRIPTION OF THE DRAWINGS

on the bactericidal activity of liquid skin cleansing formulation of the invention, both with and without antibacterial agent (e.g., Triclosan or DP300<sup>(R)</sup>). An seen, bactericidal activity of the formulation increases with lactic acid content up to about 9%. At 10% and above, bactericidal activity does not increase with increasing lactic acid content.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to liquid skin cleansing compositions comprising 1 to 99% by weight, preferably 2 to 85%, more preferably 3 to 40% of a mild surfactant system comprising one or more surfactants which alone or together have been clinically tested to be milder than soap itself as measured by zein solubilization test (soap yields 80% zein solubilized). Preferably, the mildness is such that zein solubilization is in the range 10-60% solubilization.

A number of anionic, nonionic, cationic and amphoteric surfactants may be employed in the surfactant system of the invention provided of course that the surfactant, if used alone, or surfactant mixture is milder than would be soap itself as measured by the zein solubilization test.

Among suitable anionic co-actives are the alkyl ether sulfates, acyl isethionates, alkyl ether sulfonates, sarcosinates, sulfosuccinates, taurates and combinations thereof. Among suitable amphoteric co-actives amy be included alkylbetaines, amidopropyl betaines, amidopropyl sultaines and combinations thereof.

Alkyl ether sulfates of the present invention will be of the general formula

- 10 R- $(OCH_2CH_2)_nOSO_3-M^*$  wherein R ranges from  $C_8-C_{20}$  alkyl, preferably  $C_{12}-C_{15}$  alkyl, n is an integer from 1 to 40, preferably from 2 to 9, optimally about 3, and  $M^*$  is a sodium, potassium, ammonium or triethanolammonium cation.
- 15 Typical commercial co-actives of this variety are listed in the Table below:

Trademark	Chemical Name	Physical Form	Manufacturer
Steol CS 330	Sodium Laureth Sulfate	Liquid	Stepan
Standopol ES-3	Sodium Laureth Sulfate	Liquid	Henkel
Alkasurf ES-60	Sodium Laureth Sulfate	Paste	Alkaril
Cycloryl TD	TEA Laureth Sulfate	Paste	Cyclo
Standapol 125-E	Sodium Laureth-12 Sulfate	Liquid	Henkel
Cedepal TD407MF	Sodium Trideceth Sulfate	Paste	Miranol
Standopol EA-2	Ammonium Laureth Sulfate	Liquid	Henkel

Alkyl ether sulfonates may also be employed for the present invention. Illustrative of this category is a commercial product known as Avenel S-150 commonly known as a sodium C<sub>12</sub>-C<sub>15</sub> Pareth-15 sulfonate.

Another co-active type suitable for use in the present invention is that of the sulfosuccinates. This category is best represented by the monoalkyl sulfosuccinates having the formula  $RO_2CCH_2CH(SO_3--Na^*)COO--M^*$ ; and amido-MEA sulfosuccinates of the formula  $RCONHCH_2CH_2O_2CCH_2CH(SO_3--M^*)COO--M^*$ ; wherein R ranges from  $C_8.C_{20}$  alkyl, preferably  $C_{12}-C_{15}$  alkyl and  $M^*$  is a sodium, potassium, ammonium or triethanolammonium cation. Typical commercial products representative of these co-actives are those listed in the Table below:

10

5

Trademark	Chemical Name	Physical Form	Manufacturer
Emcol 4400-1	Disodium lauryl Sulfosuccinate	Solid	Witco
Witco C5690	Disodium Cocoamido MEA Sulfosuccinate	Liquid	Witco
McIntyre Mackanate CM40F	Disodium Cocoamido MEA Sulfosuccinate	Liquid	McIntyre
Schercopol CMSNa	Disodium Cocoamido MEA Sulfosuccinate	Liquid	Scher
Emcol 4100M	Disodium Myristamido MEA Sulfosuccinate	Paste	Witco
Schercopol	Disodium Oleamido MEA	Liquid	Scher
Varsulf S13333	Disodium Ricionoleamido MEA Sulfosuccinate	Solid	Scherex

Sarcosinates may also be useful in the present invention as a co-active. This category is indicated by the general formula  $RCON(CH_3)CH_2CO_2--M^2$ , wherein R ranges from  $C_8-C_{20}$  alkyl, preferably  $C_{12}-C_{15}$  alkyl and  $M^2$  is a sodium, potassium ammonium or triethanolammonium cation. Typical commercial products representative of these co-actives are those listed in the Table below:

20

15

Trademark	Chemical Name	Physical Form	Manufacturer	
Hamposyl L-95	Sodium Lauroyl Sarcosinate	Solid	W. R. Grace	
Hamposyl TOC-30	TEA Cocoyl/Sarcosinate	Liquid	W. R. Grace	

Taurates may also be employed in the present invention as co-actives. These materials are generally identified by the formula  $RN^*(CH_3)_2CH_2CO_2-M^*$ , wherein R ranges from  $C_8-C_{20}$  alkyl, preferably  $C_{12}-C_{15}$  alkyl,  $R^1$  ranges from  $C_1-C_4$  alkyl, and  $M^*$  is a sodium, potassium, ammonium or triethanolammonium cation. Typical commercial products representative of these co-actives are those listed in the Table below:

Trademark	Chemical Name	Physical Form	Manufacturer
Igepon TC 42	Sodium Methyl Cocoyl Taurates	Paste	GAF
Igepon T-77	Sodium Methyl Oleoyl Taurate	Paste	GAF

10

Within the category of amphoterics there are three general categories suitable for the present invention. These include alkylbetaines of the formula RN\*(CH<sub>3</sub>)<sub>2</sub>CO<sub>2</sub>--M\*, amidopropyl betaines of the formula RCONHCH<sub>2</sub>CH<sub>2</sub>N\*(CH<sub>2</sub>)<sub>2</sub>CH<sub>2</sub>CO2--M\*, and amidopropyl sultaines of the formula RCONHCH<sub>2</sub>CH<sub>2</sub>N\*(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>SO<sub>3</sub>--M\* wherein R ranges from C<sub>8</sub>-C<sub>20</sub> alkyl, preferably C<sub>12</sub>-C<sub>15</sub> alkyl, and M\* is a sodium potassium, ammonium or triethanolammonium cation. Typical commercial products representative of these co-actives are found in the Table below:

Trademark	Chemical Name	Physical Form	Manufacturer
Tegobetaine F	Cocamidopropyl Betaine	Liquid	Goldschmidt
Lonzaine C	Cocamidopropyl Betaine	Liquid	Lonza
Lonzaine CS	Cocamidopropyl Hydroxysultaine	Liquid	Lonza
Lonzaine 12C	Coco-Betaine	Liquid	Lonza
Schercotaine MAB	Myristamidopropyl Betaine	Liquid	Lonza
Velvetex OLB-50	Oleyl Betaine	Paste	Henkel

Within the broad category of liquid actives, the most effective are the alkyl sulfates, alkyl ether sulfates, alkyl ether sulfonates, sulfosuccinates, and amidopropyl betaines.

5

Another preferred surfactant is an acyl isethionate having the formula:

10 
$$\parallel$$
 R - C - O - CH<sub>2</sub> - CH<sub>2</sub> - SO<sub>2</sub>M

in which R denotes a linear or branched alkyl group and M denotes an alkali metal or alkaline earth metal or an amine.

Another surfactant which may be used are the monoalkyl or dialkylphosphate surfactants.

20

25

30

35

Another mild surfactant which may be used, preferably used as primary surfactant in combination with other surfactants noted above, is sodium coco glyceryl ether sulfonate. While desirable to use because of its mildness properties, this coco AGS alone does not provide optimum lather creaminess. A sodium 90/10 coconut/tallow alkyl AGS distribution is preferred for creaminess. Salts other than the sodium salt such as TEA-, ammonium, and K-AGS and chain length distributions other than 90/10 coconut/tallow are usable at moderate levels. Also, some soap may be added to improve lather volume and speed of lathering. Certain secondary co-surfactants used in combination with AGS can also provide a creamier and more stable lather. These secondary surfactants should also be intrinsically mild. One secondary surfactant that has been found to be especially desirable is sodium lauroyl sarcosinate (trade name Hamposyl L, made by Hampshire Chemical).

The amphoteric betaines and sultaines noted above can be used as the sole surfactant, but are more preferred as a co-surfactant. Nonionics generally should not be used as the sole surfactant in this product if high foaming is desirable; however, they can be incorporated as a co-surfactant.

Nonionc and cationic surfactants which may be used include any one of those described in U.S. Patent No.

3,761,418 to Parran, Jr., hereby incorporated by reference into the subject application.

Soaps can be used at levels of about 1-10%. Soaps can be used at higher level provided that the surfactant mixture is milder than soap. The soaps may be added neat or made in situ via adding a base, e.g., NaOH; to convert free fatty acids.

Of course, as noted above, soaps should only be used as cosurfactants to the extent that the surfactant system is milder than soap alone.

Surfactant may comprise 1% to 30% by wt. of at least one anionic and 0.5% to 15% amphoteric.

25.

30

A preferred surfactant active system is one such that acyl isethionate comprises 1 to 15% by weight of the total composition, an anionic other than acyl isethionate (e.g., ammonium lauryl ether sulfate) comprises 1 to 15% by weight of the total composition and amphoteric comprises 0.5 to 15% by weight of the total composition.

#### BUFFERING COMPONENT

The second critical component of the liquid compositions of the invention is the compound or compounds which alone or together buffer the pH of the formulation under in-use condition such that the pH is from about 2.5 to 5.5, preferably 3.5 to 5.0.

By in-use is meant dilution of 1:0.5 to 1:100, 10 preferably 1:1 to 1:25 of the product in water during use.

This compound or compounds is a hydroxy carboxylic acid which lowers pH of the compositions in use to 2.5 to 5.5 and buffers at this pH.

15

20

35

5

The hydroxy carboxylic acids include any organic compound having at least one carboxylic acid group and at least one hydroxyl group. Preferably, the chain length of the acid should be C<sub>2</sub> to C<sub>18</sub>, more preferably C<sub>2</sub> to C<sub>12</sub>. Among the many acids which may be used include citric acid, lactic acid, glycolic acid, "-hydroxy C<sub>8</sub> acid, "-hydroxy C<sub>16</sub> acid, acylated citric acid and B-hydroxybutyric acid. A preferred acid is lactic acid.

In a second embodiment of the invention, the liquid skin cleansing compositions of the subject invention must contain an antibacterial agent. In this embodiment of the invention, the buffering component or compounds described above not only may provide antibacterial effect on its own, but also enhances (potentiates) the antibacterial effectiveness of the antibacterial agent.

The antibacterial agent can be present at a level of from about 0.001% to about 5%, typically from about 0.01% to about 2%, and preferably from about 0.01% to about 1.5% by

weight of the composition. The level is selected to provide the desired level of antibacterial activity and can be modified as desired. The preferred antibacterial agent is 2-hydroxy-4,2',4'-trichlorodiphenylether (DP300). Other antibacterial agents are set out below. Many antibacterial agents, known to those skilled int he art and disclosed in e.g., U.S. Patent Nos. 3,835,057 and 4,714,563, both incorporated hereinbefore by reference, may be used.

#### 10 Antimicrobials

Suitable antibacterial agents which may be used in the subject invention (i.e., in one embodiment of the invention) include:

```
2-hydroxy-4,2',4'-trichlorodiphenylether (DP300);
2,6-dimethyl-4-hydroxychlorobenzene (PCMX);
3,4,4'-trichlorocarbanilide (TCC);
3-trifluoromethyl-4,4'-dichlorocarbanilide (TFC);
```

2,2'-dihydroxy-3,3',5,5',6,6'-

20 hexachlorodiphenylmethane;

2,2'-dihydroxy-3,3',, 5,5'-tetrachlorodiphenylmethane;

2,2'-dihydroxy-3,3',dibromo-5,5'-

dichlorodiphenylmethane;

2-hydroxy-4,4'-dichlorodiphenylether;

2-hydroxy-3,5',4-tribromodiphenylether; and

1-hydroxyl-4-methyl-6-(2,4,4-trimethylpentyl)-2(1H)-pyridinone (Octopirox).

Other suitable antimicrobials include:

30

25

Benzalkonium chloride;
Benzethonium chloride;
Carbolic acid;
Cloflucarbon (Irgasan CF3;4,4'-dichloro-3-

35 (trifluoromethyl)carbanilide);

```
Chlorhexidine (CHX; 1,6-di(4'-chlorophenyl-diguanido)hexane);
Cresylic acid;
Hexetidine (5-amino-1,3-bid(2-ethylhexyl)-5-methylhexahydropyrimidine);
Iodophors;
Methylbenzethonium chloride;
Povidone-iodine;
Tetramethylthiuram disulfide (TMTD; Thiram);
Tribrominated salicylanilide.
```

In addition to a mild surfactant compound or compounds, the pH buffering compound or compounds, water and optionally (or as required in one embodiment), antimicrobial agent, the liquid skin cleansing compositions may contain optionals as described below.

Each of the above components can be incorporated in an aqueous vehicle which may, in addition, include such materials as organic solvents, such as ethanol, thickeners, 20 such as carboxymethylcellulose, magnesium aluminum silicate, hydroxyethylcellulose, methylcellulose or carbopols; perfumes; sequestering agents, such as tetrasodium ethylenediaminetetraacetate (EDTA), EHDP or mixtures in an 25 amount of 0.01 to 1%, preferably 0.01 to 0.05%; and coloring agents, opacifiers and perlizers such as zinc stearate, magnesium stearate, TiO,, EGMS (ethylene glycol monostearate) or Lytron 621 (Styrene/Acrylate copolymer); all of which are useful in enhancing the appearance or cosmetic properties of 30 the product.

The following preservatives may also be used in the liquid skin cleansers of the invention:

#### LIQUID SKIN CLEANSER PRESERVATIVES

PRESERVATIVE	CHEMICAL NAME
Bronopol	2-Bromo-2-nitropropane-1,3,diol
Dowicil 200	cis Isomer of 1-(3-chloroally1)-3,5,5-triaza-1- azoniadamantane-chloride OR Quaternium 15
Glycacil	3-Iodo-2-propynyl butyl carbamate
Glydant XL 1000	DMDM Hydantoin OR dimethyloldimethylhydantoin
Glydant Plus	DMDM Hydantoin and 3-iodo-2-propynyl butyl carbamate
Formaldehyde	Formaldehyde
Germall II	N-(Hydroxymethyl)-N-(1,3-dihydroxymethyl-2,5-dioxo-4- imidazolidinyl)-N'-(hydroxymethyl) urea OR Diazolidinyl urea
Germall 115	N,N'-methylene-bis-[N'-1-(hydroxymethyl)-2,,5-dioxo-4-imidazolidinyl]urea OR imidazolidinyl urea
Glutaraldehyde	Glutaraldehyde
Kathano CG	Mixture of 5-chloro-2-methyl-4-isothiazoline-3-one- and 2-methyl-4-isothiazoline-3-one OR Mixture of methyl, chloromethyl isothiazolinone, and methyl isothiazolinone
Parabens	Methyl Paraben, and Ethyl Paraben, and Propyl Paraben and Butyl Paraben OR those esters of p-hydroxybenzoic acid
Phenoxyethanol	2-Phenoxyethanol
Salicylic Acid	Salicylic Acid OR o-Hydroxybenzoic acid
Sorbic Acid	Sorbic Acid, Potassium Sorbate

Coconut acyl mono- or diethanol amides as suds boosters, and strongly ionizing salts such as sodium chloride and sodium sulfate may be used to advantage.

Antioxidants such as, for example butylated hydroxytoluene (BHT) may be used advantageously in amounts of about 0.01% or higher if appropriate.

Cationic conditioners which may be used include Quatrisoft LM-200 (Polyquaternium-24); polyethylene glycols such as

5 Polyox WSR-205 PEG 14M,

WSR-N-60K PEG 45M, or

WSR-N-750 PEG 7M; and

Merquat Plus 3330 - Polyquaternium 39.

Thickeners which may be used include Americall Polymer HM 1500 (Nonoxynyl Hydroethyl Cellulose); Glucan DOE 120 (PEG 120 Methyl Glucose Dioleate).

Unless stated otherwise, the percentages int he specification, examples and claims are percentages by weight.

Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this description indicating amounts of material (or conditions of reaction and/or use) are to be understood as modified by the word "about".

The following examples are intended for illustrative purposes only and should not be construed to limit the invention in any way.

#### EXAMPLES

An <u>In vitro</u> Bactericidal Kill Test is used to measure antimicrobial activity in the examples which follow.

Methodology for the test is set forth below:

WO 98/37866 PCT/EP98/00155

- 18 -

#### IN VITRO BACTERICIDAL KILL TEST

An <u>in vitro</u> bactericidal test was used to determine the antibacterial effect of products on <u>Staphylococcus aureus</u>
ATCC #6538 during a short contact time. One milliliter (about 10° cells) of bacteria was exposed for one minute to a one-percent solution of liquid skin cleansing composition. The sample was added to additional water, mixed, and further diluted in 0.1% peptone. Duplicate samples of appropriate dilutions were plated on neutralizing media. In addition, the bacterial culture was plated to determine the actual number of organisms inoculated. the plates were incubated at 34°C for 48 hours and enumerated. The CFR/ml (colony forming units per milliliter) from dilutions with plate counts int he range of 30-300 were averaged together to produce the final CFU/ml.

The results may be expressed as the log of the CFU/ml. The culture control indicates the actual number of bacteria inoculated while the water control reflects the number of organisms recovered int he absence of product. The lower the number, the more effective the solution was in killing the bacteria.

In this assay, a sampling error of approximately 0.5 log is likely, therefore differences of 0.5 log between products may not be significant. As a result, the data should be viewed in terms of trends rather than as absolute numbers.

30

5

15

20

#### Example 1

Applicants carried out an experiment showing that lactic acid concentration on the bacteriocidal activity of liquid skin cleansing formulation. As seen in Figure 1, the

bacteriocidal activity of the formulation increases with lactic acid content up to about 9%. At 10% and above, bactericidal activity does not incease with increasing lactic acid content.

5

INGREDIENT	% BY WEIGHT
Anionic (Acyl Isethionate)	1 to 15%
Anionic Other than Acyl Isethionate (e.g., SLES)*	1 to 15%
Amphoteric Surfactant **	5 to 15%
pH Buffering (Lactic Acid)	1 to 5%
Sequestrant (EDTA or EHDP)	0.01 to 0.1%
Moisturizer (e.g., Cationic Polymer)	0.05 to 3.0%
Additives (e.g., Dyes, Perfumes)	0 to 10%
Water	Balance

- \* SLES sodium lauryl ether sulfate
- \*\* Cocoamidopropyl betaine

#### 10 Example 2

The compound or compounds of the invention may also be used in the following formulations.

FORMULATION 1			
COMPONENT	% BY WEIGHT		
Sodium Isethionate	3-5%		
Sodium Alkene Benzene Sulfonate	1-3%		
Sodium Laureth Sulfate	3-5%		
Sodium Cocoyl Isethionate	8-12%		
Sodium Tallow/Coconut Soap	1-3%		
Preservative (e.g., Methylparaben)	.15%		
Sequestrants	.0105%		
Fatty Acid (e.g., Stearic Acid)	7-10%		
Sulfosuccinate	3-5%		
Water plus minors	to balance		

FORMULATION 2		
COMPONENT	% BY WEIGHT	
Sodium Cocoyl Isethionate	5-8%	
Cocamidopropyl Betaine	5-8%	
Sulfosuccinate	2-5%	
Fatty Acid	6-9%	
Sodium Isethionate	1-3%	
Silicone Emulsion	3-7%	
Sequestrant	.0105%	
Water plus minors	to balance	

#### **CLAIMS**

- 1. A single phase skin cleansing composition comprising:
- 5 (1) 1% to 99% by weight of a surfactant system comprising:
  - (a) 1% to 30% by wt. of at least one anionic surfactant; and
  - (b) 0.5% to 15% of an amphoteric surfactant.

10


15

- (2) 0.5% to 9% by wt. of a hydroxy carboxylic acid compound or compounds which buffers pH of the composition such that pH is less than 5 upon dilution with water at ranges of 1:0.5 to 1:100 dilution; and
- (3) 1% to 99% by wt. water.
- A skin cleansing composition as claimed in claim 1
   further comprising 0.001% to about 5% by weight of an antibacterial agent.
- A composition as claimed in either claim 1 or claim 2, wherein the surfactant system is 2-85% by wt. of the
   composition.
  - 4. A composition as claimed in any preceding claim, wherein the surfactant systems is 3-40% by weight of the composition.

30

- 5. A composition as claimed in any preceding claim, wherein pH is from about 2.5 to less than 5.0.
- 6. A composition as claimed in any preceding claim, 35 wherein pH is from about 3.0 to less than 5.0.

- 7. A composition as claimed in any preceding claim, wherein the hydroxy carboxylic acid is lactic acid.
- 5 8. A composition as claimed in any preceding claim, wherein the surfactant system comprises 1 to 15% by wt. acyl isethionate.



#### INTERNATIONAL SEARCH REPORT

In tional Application No PCT/FP 98/00155

<del></del>			101/61 30/00133	
A. CLASS IPC 6	SIFICATION OF SUBJECT MATTER A61K7/50 A61K7/48			-
According	to International Patent Classification (IPC) or to both national di	assification and IPC	,	
	S SEARCHED			
Minimum d IPC 6	ocumentation searched (classification system followed by class A61K	sification symbols)		
Documenta	ation searched other than minimum documentation to the extent	that such documents are inclu	ded in the fields searched	
Electronic	data base consulted during the international search (name of d	ata base and, where practical,	search terms used)	
C. DOCUM	MENTS CONSIDERED TO BE RELEVANT	<del></del>		
Category *		he relevant passages	Relevant to claim N	
			1 10/0 value to carried by	J.
X	WO 95 32705 A (UNILEVER PLC ; (NL)) 7 December 1995 see the whole document	UNILEVER NV	1-8	
X	GB 2 288 811 A (PROCTER & GAME November 1995 see the whole document			
X	WO 94 17166 A (PROCTER & GAMBI MICHEL JOSEPH (GB); LEAHY CHRI DAVI) 4 August 1994 see page 1-7		1-6	
	see page 14, line 24-30 see page 16, line 18-19 see page 16, line 30-31 see examples 1-7			
		-/ <del></del>		
X Furt	ther documents are listed in the continuation of box C.	X Patent family m	embers are listed in annex.	
"A" docume consice "E" earlier of filling of which citation "O" docume other of docume other of docume other of docume other of the citation o	ategories of cited documents:  ent defining the general state of the art which is not dered to be of particular relevance document but published on or after the international date ent which may throw doubts on priority claim(s) or is cited to establish the publicationdate of another in or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or means ent published prior to the international filing date but han the priority date claimed	or priority date and cited to understand invention  "X" document of particul cannot be consider involve an inventive "Y" document of particul cannot be consider document of cannot be considered to comment is combile.	shed after the international filing date not in conflict with the application but the principle or theory underlying the ar relevance; the claimed invention ed novel or cannot be considered to step when the document is taken alone ar relevance; the claimed invention ad to involve an inventive step when the led with one or more other such documention being obvious to a person skilled	
<u> </u>	actual completion of theinternational search	<del></del>	e international search report	
	June 1998	16/06/19	98	
Name and r	mailing address of the ISA  European Patent Office, P.B. 5818 Patentlaan 2  NL - 2280 HV Rijswijk  Tel. (+31-70) 340-2040, Tx. 31 651 epo nt.	Authorized officer	onzalez M	

#### INTERNATIONAL SEARCH REPORT

Int tional Application No PCT/EP 98/00155

C.(Continue	PCT/EP 98/00155  nuation) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the relevant passages  Relevant to daim No.		
Category '			
<b>(</b>	EP 0 662 316 A (OREAL) 12 July 1995 see the whole document	1,3-7 1-8	
<b>'</b>	WO 96 21426 A (PROCTER & GAMBLE) 18 July 1996 see page 1-10 see page 17, line 29	1-8	
	·		
	• •	-	
	į.		
	·		
	·		

#### INTERNATIONAL SEARCH REPORT

information on patent family members

PCT/EP 98/00155

	tent document in search report	t	Publication date	Patent family member(s)	Publication date
WO	9532705	A	07-12-1995	US 5681802 A AU 2735595 A BR 9507819 A CA 2186011 A	28-10-1997 21-12-1995 16-09-1997 07-12-1995
				CZ 9603500 A EP 0762868 A HU 76537 A	14-05-1997 19-03-1997 29-09-1997
				JP 10500962 T PL 317427 A SK 152596 A	27-01-1998 14-04-1997 04-06-1997
GB	2288811	A	01-11-1995	NONE	
WO	9417166	A	04-08-1994	CN 1116858 A EP 0689581 A JP 8505875 T	14-02-1996 03-01-1996 25-06-1996
EP	0662316	A	12-07-1995	FR 2714826 A AT 145546 T CA 2138244 A DE 69400996 D DE 69400996 T ES 2097624 T HU 71738 A JP 2603810 B JP 8034710 A PL 306578 A	13-07-1995 15-12-1996 16-06-1994 09-01-1997 27-03-1997 01-04-1997 29-01-1996 23-04-1997 06-02-1996 24-07-1995
WO	9621426	Α	18-07-1996	EP 0802786 A	29-10-1997